# **Evolution of an atypical de-epoxidase for photoprotection in the green lineage**

Zhirong Li<sup>1,2</sup>, Graham Peers<sup>3</sup>, Rachel M. Dent<sup>1</sup>, Yong Bai<sup>1</sup>, Scarlett Y. Yang<sup>1†</sup>, Wiebke Apel<sup>1†</sup>, Lauriebeth Leonelli<sup>1</sup> and Krishna K. Niyogi<sup>1,2\*</sup>

Plants, algae and cyanobacteria need to regulate photosynthetic light harvesting in response to the constantly changing light environment. Rapid adjustments are required to maintain fitness because of a trade-off between efficient solar energy conversion and photoprotection. The xanthophyll cycle, in which the carotenoid pigment violaxanthin is reversibly converted into zeaxanthin, is ubiquitous among green algae and plants and is necessary for the regulation of light harvesting, protection from oxidative stress and adaptation to different light conditions<sup>1,2</sup>. Violaxanthin de-epoxidase (VDE) is the key enzyme responsible for zeaxanthin synthesis from violaxanthin under excess light. Here we show that the Chlorophycean VDE (CVDE) gene from the model green alga Chlamydomonas reinhardtii encodes an atypical VDE. This protein is not homologous to the VDE found in plants and is instead related to a lycopene cyclase from photosynthetic bacteria<sup>3</sup>. Unlike the plant-type VDE that is located in the thylakoid lumen, the Chlamydomonas CVDE protein is located on the stromal side of the thylakoid membrane. Phylogenetic analysis suggests that CVDE evolved from an ancient de-epoxidase that was present in the common ancestor of green algae and plants, providing evidence of unexpected diversity in photoprotection in the green lineage.

Photosynthetic organisms are subjected to a large dynamic range of light intensities, which can vary rapidly because of canopy shading, passing clouds or sunflecks, as well as on a daily or seasonal basis. To allow optimal photosynthesis at low light intensities and to avoid photo-oxidative damage because of the formation of reactive oxygen species (ROS) under excess light, photosynthetic organisms have evolved the ability to regulate light harvesting. Under excess light, photosynthetic light harvesting is regulated by non-photochemical quenching (NPQ) mechanisms that are responsible for dissipating excess absorbed light as heat<sup>4-7</sup>. The major and most intensively investigated component of NPQ is called qE, which is turned on and off on the time scale of seconds to minutes. qE depends on acidification of the thylakoid lumen on formation of high  $\Delta pH$  across the thylakoid membrane in excess light<sup>8</sup>. In plants, this results in two important changes that facilitate qE: conformational changes of light-harvesting complex proteins by protonation and the activation of a lumen-localized violaxanthin (Vio) de-epoxidase (VDE) enzyme. VDE catalyses the conversion of Vio to zeaxanthin (Zea) via the intermediate antheraxanthin (Anthera). Zea and Anthera (xanthophylls with a de-epoxidized 3-hydroxy  $\beta$ -ring end group) are the major xanthophyll pigments that are involved in qE in plants. Zea epoxidase converts Zea back

to Vio in limiting light. Together, these light intensity-dependent interconversions are known as the xanthophyll cycle (Fig. 1a). Xanthophyll de-epoxidation occurs in almost all photosynthetic eukaryotes, although it contributes to qE and other NPQ mechanisms to different extents in different organisms<sup>9-11</sup>. In green algae and plants, Zea also plays important roles in photoprotection as an antioxidant that directly quenches singlet oxygen and triplet chlorophyll species<sup>12-14</sup>.

Mutants defective in the xanthophyll cycle and qE have been identified in the unicellular green alga *Chlamydomonas reinhardtii* and the model plant *Arabidopsis thaliana*<sup>15,16</sup>. The *npq1* mutants are defective in VDE activity and are unable to convert Vio to Anthera and Zea in high light (Fig. 1a,d). Although the *Arabidopsis npq1* mutant was shown to affect the *VDE* gene<sup>16</sup>, the molecular basis of the *Chlamydomonas npq1* mutant has been mysterious, because the *Chlamydomonas* genome lacks an obvious orthologue of the *VDE* gene found in plants and other algae. Furthermore, VDE activity is not inhibited by dithiothreitol (DTT) in *Chlamydomonas* cells<sup>11</sup>, unlike in plants, indicating that *Chlamydomonas* is most likely to employ a novel type of VDE.

The Chlamydomonas npq1 mutation had been previously mapped to linkage group IV17. By fine mapping, we localized the npq1 mutation to a small region containing 13 gene models as candidate genes. One of these gene models (Cre04.g221550) encodes a putative flavin adenine dinucleotide (FAD)-dependent oxidoreductase with a predicted chloroplast transit peptide. Genomic polymerase chain reaction (PCR) analysis showed that there was a 164 bp deletion in the *npq1* allele (Fig. 1b and Supplementary Fig. 1) of this gene. Introducing a Cre04.g221550 genomic clone into the npq1 mutant strain restored Zea synthesis in high light (Fig. 1d). Interestingly, some rescued lines accumulated higher levels of Zea than the wild type (Fig. 1d), which correlated with higher accumulation of the protein encoded by Cre04. g221550 (Supplementary Fig. 2). From the results of these experiments, it is clear that the Zea deficiency of npq1 is caused by the loss of Cre04.g221550 function.

To determine if Cre04.g221550 actually encodes a protein with VDE activity, we tested if this gene could complement the *Arabidopsis npq1* mutation (here called *vde1*), which is known to disrupt the endogenous plant-type *VDE* gene<sup>16</sup>. To ensure proper expression and chloroplast targeting of the Cre04.g221550 protein, we codon optimized the *Cre04.g221550* gene sequence for *Arabidopsis*, either with a sequence encoding its native, amino (N)-terminal chloroplast transit peptide or the chloroplast transit peptide from the *Arabidopsis* PsbS protein, and with or without a

<sup>&</sup>lt;sup>1</sup>Department of Plant and Microbial Biology, Howard Hughes Medical Institute, University of California, Berkeley, California 94720-3102, USA. <sup>2</sup>Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA. <sup>3</sup>Department of Biology, Colorado State University, Fort Collins, Colorado 80523-1878, USA. <sup>†</sup>Present addresses: Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA (S.Y.Y.). Institute for Biology, Experimental Biophysics, Humboldt-Universität zu Berlin, 10115 Berlin, Germany (W.A.). \*e-mail: niyogi@berkeley.edu



**Figure 1** | Molecular analysis and complementation of *npq1* mutation in *Chlamydomonas*. **a**, Xanthophyll cycle reactions. The de-epoxidation of violaxanthin to zeaxanthin via antheraxanthin is defective in the *npq1* mutant. **b**, Schematic showing the *Cre04.g221550* (*CrCVDE*) gene model and the 164 bp deletion in the *npq1* mutant allele. **c**, Phylogenetic analysis of CVDE and CruP proteins. Syn, *Synechococcus* sp. strain PCC7002; Phys, *Physcomitrella patens*; At, *Arabidopsis thaliana*; Ostta, *Ostreococcus tauri*; Cr, *Chlamydomonas reinhardtii*; Vocar, *Volvox carteri*. Scale bar indicates the substitution rate of nucleotides per site. Branch support values were generated by the approximate likelihood-ratio test. **d**, High-performance liquid chromatography (HPLC) phenotype of wild type, *npq1* and two independent complemented lines. Arrows denote the Zea peak resulting from CVDE activity.

carboxy (C)-terminal FLAG epitope tag (Fig. 2a). Arabidopsis vde1 lines expressing each of the four versions of *Cre04.g221550* displayed excess light-induced Zea synthesis and NPQ phenotypes similar to wild-type plants (Fig. 2b,c), showing that the *Cre04. g221550* gene indeed encodes a functional, evolutionarily distinct VDE enzyme. Based on the presence of homologues of *Cre04. g221550* in sequenced green algae of the class Chlorophyceae, we designate this gene *CVDE* to distinguish it from the plant-type *VDE* gene.

Plant-type VDE is localized in the thylakoid lumen and associates with the thylakoid membrane, where it catalyses the de-epoxidation reaction on membrane-associated Vio. We used lines of both the *Chlamydomonas npq1* mutant and the *Arabidopsis vde1* mutant complemented with a C-terminal FLAG-tagged version of the *Chlamydomonas* CVDE (CrCVDE) protein to determine its localization. The functional C-terminal tagging demonstrated that this modification does not impair CrCVDE enzyme activity (Fig. 2b,c). Using either a polyclonal antibody raised against an N-terminal 15 amino acid peptide of mature CrCVDE or a commercial antibody raised against the FLAG epitope, we detected the CrCVDE protein at a molecular mass of 87 kDa (Fig. 3), which is the predicted size of the mature protein after cleavage of the chloroplast transit peptide. As expected, the CrCVDE protein is associated with the thylakoid membrane in

both Chlamydomonas and Arabidopsis (Fig. 3a-c). To determine the topology of CrCVDE, we performed a limited proteolysis experiment with isolated thylakoid membranes from both Chlamydomonas- and Arabidopsis-complemented lines. Thermolysin treatment resulted in complete cleavage of the CrCVDE protein, even more rapidly than the cleavage of the stroma-exposed PsaD subunit of photosystem I, which was quickly digested to a thermolysin-resistant fragment (Fig. 3b,c). In contrast, the PsbO subunit of photosystem II, located in the thylakoid lumen, was completely resistant to thermolysin unless the membrane was solubilized with detergent (Fig. 3b,c). In Arabidopsis, the lumen-localized planttype VDE protein (in the vde1 mutant complemented with a FLAG-tagged version of the Arabidopsis VDE gene) was not affected unless the membrane was solubilized with detergent (Fig. 3c). These results strongly suggest that the epitope-tagged CrCVDE protein is located on the stromal side of the thylakoid membrane when expressed in either Chlamydomonas or in Arabidopsis (Fig. 3d), which differs from the plant-type VDE that is located in the thylakoid lumen (Fig. 3c). The stroma-exposed location of CrCVDE was further supported by the presence of an FAD-binding domain in the mature CrCVDE protein (FAD is present in the stroma but not the thylakoid lumen). Salt wash assays indicated that CrCVDE is peripherally associated with the membrane and could be extracted by NaSCN (Supplementary Fig. 3).



**Figure 2 | Functional complementation of** *Arabidopsis vde1* **mutant by expression of the** *Chlamydomonas* **CVDE protein. a**, Constructs used for transformation of the *Arabidopsis vde1* mutant and their ability to complement the zeaxanthin accumulation and NPQ phenotypes. '+' indicates successful rescue of the phenotype. **b**, HPLC phenotypes of wild type, *vde1* mutant and two complemented lines. The arrows denote the Zea peak resulting from CVDE activity (or plant-type VDE activity in the wild type). c, NPQ induction and relaxation of wild type, *vde1* mutant and two independent complemented lines. The white bar above the graph indicates illumination with 1,250 µmol photons  $m^{-2} s^{-1}$ ; the black bar indicates darkness (with only very weak measuring light). Error bars represent standard deviation (*n* = 5).

The in vivo substrate of VDE, Vio, is free in the membrane lipid phase rather than bound to pigment proteins<sup>2,18</sup>. Therefore, one possible explanation of functional replacement of plant-type VDE in Arabidopsis by CrCVDE is that substrate Vio molecules are accessible to enzymes on either side of the thylakoid membrane (in the thylakoid lumen or in the stroma of the chloroplast). Indeed, addition of partially purified plant-type VDE from spinach to the stromal side of thylakoids isolated from the Arabidopsis vde1 mutant rescued the mutant phenotype in vitro<sup>19</sup>. Similar to planttype VDE, the CVDE activity of intact Chlamydomonas cells was inhibited by the uncoupler nigericin (Supplementary Fig. 4), indicating that the activation of this stromal enzyme also requires the build-up of a large pH gradient in excess light. Plant-type VDE requires ascorbate to catalyse the de-epoxidation reaction, but at this time it is not clear what other substrates are required for CVDE activity.

The evolutionary origins of plant-type VDE and CVDE are clearly distinct. CVDE is a homologue of CruP and CruA (Fig. 1c

and Supplementary Fig. 5). CruA is known to be involved in bacterial carotenoid biosynthesis as a lycopene cyclase<sup>3</sup>, whereas CruP is a paralogue of CruA. We note that the proposed carotenoid cyclase<sup>20</sup> and de-epoxidase reaction mechanisms are similar (Supplementary Fig. 6), suggesting that a de-epoxidase enzyme could evolve from a cyclase. Our demonstration that CrCVDE has VDE activity suggests that its paralogue CruP, which is widely distributed in oxygenic photosynthetic organisms<sup>21</sup>, might also be a de-epoxidase. Based on the observation that cruP mutants or overexpressors of Arabidopsis accumulate more or less β-carotene-5,6epoxide (an oxidized derivative of  $\beta$ -carotene), respectively, when challenged by stress<sup>21</sup>, we hypothesize that CruP is a  $\beta$ -carotene-5,6-epoxide de-epoxidase. CVDE and CruP homologues are present in Chlamydomonas and its multicellular relative Volvox carteri, but only CruP homologues can be found in Ostreococcus tauri, Arabidopsis thaliana and Physcomitrella patens. Phylogenetic analysis strongly suggests that CVDE evolved by duplication of CruP in the ancestor of green algae and plants and that CVDE



**Figure 3 | Subcellular localization of CrCVDE proteins expressed in Chlamydomonas and Arabidopsis. a**, Immunoblot analysis of total cell (T), membrane (M) and soluble (S) fractions of *Chlamydomonas* strains. The FLAG-tagged CrCVDE protein is detected in the membrane fraction and not in the soluble fraction in two independent transformants. Subcellular markers: D2 for membrane fraction and RbcL for soluble fraction. b, Protease protection assay of isolated intact thylakoids from *Chlamydomonas* complemented lines. Isolated thylakoids were treated with thermolysin in the presence and absence of Triton X-100. Aliquots were removed at the specified times, samples were separated by SDS-PAGE and immunodetection was performed with specified antibodies. Thermolysin-resistant Atpβ was used as a loading control. The FLAG-tagged CrCVDE protein was probed with both the N-terminal epitope antibody and the C-terminal FLAG antibody. Subcellular markers: PsaD for stroma-exposed membrane protein; PsbO for thylakoid lumen. **c**, Immunoblot analysis and protease protection assay of the FLAG-tagged CrCVDE protein expressed in the *Arabidopsis vde1* mutant. Left panel, the CrCVDE protein is detected in the thylakoid lumen and RbcL for stroma. Right panel, protease protection assay of isolated thylakoids from *Arabidopsis* complemented lines. RbcL was not present in the thylakoid fraction. Lower panel, the location of the plant-type VDE in the thylakoid lumen was confirmed by analysis of a transformant expressing the FLAG-tagged *Arabidopsis* VDE protein in the *vde1* mutant. The migration of the VDE protein in the chloroplast fraction is altered by the comigration of a protein that is absent from the thylakoid fraction. **d**, Proposed topology of CrCVDE in both *Chlamydomonas* and *Arabidopsis*.

has been selectively lost in some clades of the Viridiplantae (Fig. 1c). This might explain some previous observations of DTT-resistant VDE activity in green algae<sup>9–11</sup>; however, the limited number of genomes sequenced within this clade prohibits any further speculation about the distribution or origin of CVDE-related xanthophyll cycling.

The evolutionary history of algae (and plants) is complicated by endosymbiosis and horizontal gene transfer events. We showed that a novel de-epoxidase from a green algal group is functional in a land plant, despite their evolutionary separation by more than 700 million years<sup>22</sup>. Therefore, it may be possible to mix and match the regulatory components of light harvesting from different clades of photosynthetic organisms to effectively tune photosynthetic efficiency and increase photosynthetic productivity.

### Methods

Genetic mapping and PCR analysis. The fine mapping of the npq1 mutation was done by scoring PCR-based markers on selected tetrad mutant progeny derived from a cross between npq1 (137c strain background) and the polymorphic wild-type strain S1D2 (CC-2090). Markers were designed based on information in

Kathir *et al.*<sup>23</sup> and the marker list from David Stern available at www.chlamy.org. To identify the mutation in the *CVDE* gene, genomic DNA PCR was performed with a series of primer pairs that collectively span the entire gene, and the PCR products were sequenced for comparison between the wild type and the *npq1* mutant. The primers that resulted in different length products between wild type and *npq1* were RMD345 (5'-CTTGGCGGAAGCAGAGTATGGC-3') and RMD346 (5'-CGGCCTCCCTTCATCCCTCCCAC-3').

**Phylogenetic analysis.** CVDE homologues and CruP homologues were identified by searching using BlastP and tBlastN against the sequenced proteome and genome database, respectively, with an e-value cutoff of  $1 \times 10^{-90}$ . The potential chloroplast transit peptides for CVDE homologues or CruP homologues were predicted by aligning respective homologues from organisms with or without chloroplasts using the Clustal Omega program (version1.2.1; http://www.ebi.ac.uk/Tools/msa/clustalo/). The predicted mature proteins were aligned using Clustal Omega and BoxShade (version 3.21; www.ch.embnet.org/software/BOX\_form.html). The phylogenetic tree was constructed at Phylogeny.fr (http://phylogeny.lirmm.fr/phylo\_cgi/ advanced.cgi) with Gblocks for alignment curation, PhyML for construction of the Phylogenetic tree.

**Complementation of** *Chlamydomonas npq1* **mutant.** For complementation of *npq1*, an 11.5-kb *Eco*RV/*Not*I fragment of BAC clone 33B9 containing the *CVDE* gene was subcloned into the pBC1 vector<sup>24</sup> to generate pCVDEg. For

### NATURE PLANTS DOI: 10.1038/NPLANTS.2016.140

# LETTERS

complementation of *npq1* with a C-terminal FLAG-tagged version of the CVDE protein, the 1.4 kb *Sbfl/Bgl*II fragment of pCVDEg containing the 3'-terminus of the *CVDE* gene was subcloned into the pUC19–BglII vector to generate pUC19–BglII-pCVDE. The 0.4 kb *Ncol/Bg*III fragment of pUC19–BglII-pCVDE was then replaced by a synthesized version (Integrated DNA Technologies, Inc.), which contains a C-terminal FLAG-tag linked with the CVDE protein through two glycines to generate plasmid pUC19–BglII-pCVDE–FLAG. The 1.4 kb *Sbfl/Bg*III fragment of pUC19–BglII-pCVDE–FLAG. The 1.4 kb *Sbfl/Bg*III fragment of pUC19–BglII-pCVDE–FLAG was then ligated into pCVDEg double-digested with the same enzymes to generate pCVDEg–FLAG. Both pCVDEg and pCVDEg–FLAG were separately transformed into the *npq1* mutant using the glass bead method as described previously<sup>25</sup>. The positive transformants were selected on paromomycin and then screened for zeaxanthin accumulation after high light exposure by high-performance liquid chromatography (HPLC) as previously described<sup>26</sup>.

Complementation of Arabidopsis vde1 mutation by Chlamydomonas CVDE. The predicted protein sequences of Chlamydomonas CVDE were retrieved from both Phytozome at http://www.phytozome.net (protein ID: Cre04.g221550.t1.2) and the Joint Genome Institute at http://genome.jgi-psf.org/Chlre4/Chlre4.home.html (protein ID: 522089). The predicted CVDE protein sequences were confirmed by comparing against each other and against the cDNA consensus obtained from UCSC/UCLA genome browser at http://genomes.mcdb.ucla.edu. The CDS of the CrCVDEAt gene was then codon optimized and synthesized for Arabidopsis nuclear/ cytoplasmic expression (GenScript). The synthetic CrCVDEAt gene was subcloned into the Gateway vector pDONR221, and a FLAG tag was added right before the stop codon by 'Round-the-horn' site-directed mutagenesis (http://openwetware.org/ wiki/%27Round-the-horn\_site-directed\_mutagenesis). Sequence encoding the Arabidopsis PSBS transit peptide (first 54 amino acids) was amplified to replace the predicted native CrCVDE transit peptide (first 56 amino acids) in versions of each construct using gene SOEing<sup>27</sup>. The CrCVDEAt gene and the FLAG-tagged CrCVDEAt gene were subcloned into the pEarleyGate100 vector<sup>28</sup> and transformed into the Arabidopsis vde1 mutant<sup>16</sup> using the floral dip method<sup>29</sup>. As a positive control, a vector containing a FLAG-tagged version of the Arabidopsis VDE1 gene<sup>30</sup> was also transformed. The transformants were selected on Murashige and Skoog plates containing 20 μg ml<sup>-1</sup> glufosinate ammonium, screened for NPQ capacity with the IMAGING-PAM M-series (Heinz Walz), measured for NPQ induction with an FMS2 fluorometer (Hansatech Instruments) as previously described<sup>31</sup>, and assayed for the accumulation of zeaxanthin after high light exposure by HPLC as described<sup>26</sup>.

**Chlamydomonas cell fractionation.** Chlamydomonas cells were grown photoheterotrophically in TAP medium<sup>32</sup> to medium logarithmic phase (approximately  $5 \times 10^6$  cells ml<sup>-1</sup>) and harvested by centrifugation at 3,000g for 5 min. Cells were resuspended in PBS buffer to a density of  $2 \times 10^8$  cells ml<sup>-1</sup> and broken by FastPrep-24 (MP Biomedicals) with lysing matrix J at a speed of 4.0 m s<sup>-1</sup> for 40 s. Total membrane and total supernatant were separated by centrifugation at 20,000g and 4 °C for 10 min. Total membranes were washed three times before being resuspended with 1× PBS buffer containing 100  $\mu$ M phenylmethylsulphonyl fluoride (PMSF). Samples were then subjected to immunoblot analysis as described below.

Chlamydonas and Arabidopsis thylakoid isolation. Chlamydomonas thylakoids were isolated by a modification of the flotation procedure described previously<sup>33</sup> The Chlamydomonas cells were grown in 400 ml of TAP under low light and harvested at the mid-logarithmic growth phase. The cell pellet was resuspended in 20 ml of 25 mM HEPES (pH 7.5), 0.3 M sucrose, 10 mM CaCl<sub>2</sub> and 10 mM MgCl<sub>2</sub> with protease inhibitors. The cells were broken by passing the resuspended cells through a chilled French pressure cell, and the homogenate was centrifuged at 18,000 rpm for 10 min. The supernatant was discarded and the pellet was gently resuspended with a paintbrush in 5 ml of 5 mM HEPES (pH 7.5), 1.8 M sucrose, 10 mM CaCl<sub>2</sub> and 10 mM MgCl<sub>2</sub>. The resuspension was carefully transferred into a clear tube for the SW41 rotor and topped with 6 ml of 5 mM HEPES (pH 7.5), 0.5 M sucrose, 10 mM CaCl2 and 10 mM MgCl2. The tubes were centrifuged at 38,000 rpm (SW41, 4 °C) for 1 hour. The membrane layer at the interface of two solutions was carefully transferred to a 1.5 ml Eppendorf tube containing 1 ml of 25 mM HEPES (pH 7.5), 0.3 M sucrose, 10 mM CaCl<sub>2</sub> and 10 mM MgCl<sub>2</sub>.

Fresh *Arabidopsis* rosette leaves were harvested from 4-week-old plants grown in controlled conditions of 14 h light, 22 °C/10 h dark, 23 °C, with a light intensity of 150  $\mu$ mol photons m<sup>-2</sup> s<sup>-1</sup> and stored on ice. The *Arabidopsis* thylakoids were isolated from the leaves as previously described<sup>34</sup>.

**Protease protection assay.** Thylakoids were resuspended in 0.3 M sorbitol, 2.5 mM EDTA, 5 mM MgCl<sub>2</sub>, 0.5% (w/v) BSA, 20 mM HEPES (pH 7.6) at 0.3 nmol of chlorophyll *a* per ml. The reaction was started by the addition of thermolysin (EMD Millipore) at a final concentration of 20  $\mu$ g ml<sup>-1</sup> to 400  $\mu$ l of thylakoids preparation. The reaction was stopped by transferring 60  $\mu$ l to a tube containing 6  $\mu$ l of 500 mM EDTA at six different time points: 0, 0.5, 2, 5, 10 and 20 min. The tubes were vortexed immediately, and 66  $\mu$ l of 2× sample buffer was added.

CVDE antibody generation and immunoblot analysis. The polyclonal antibody recognizing CrCVDE was generated in rabbits against an epitope located near the N terminus of the protein sequence of CrCVDE (CLRNQKHEPEKKGPK), and the resulting crude serum was affinity purified (ProSci Inc.). Polyclonal antibodies against D2, PsbO, PsaD and RbcL were obtained from Agrisera (Sweden) and FLAG antibody was from ThermoFisher Scientific. Protein samples were solubilized with 2× solubilization buffer (500 mM Tris-HCl (pH 6.8), 7% SDS, 20% glycerol (v/v), 2 M urea and 10%  $\beta$ -mercaptoethanol (v/v)) by pipetting up and down several times before incubation at room temperature for 30 min. For immunoblot analysis of CVDE, protein samples were separated with NuPAGE Novex 3-8% Tris-Acetate mini gels (Life Technologies). For immunoblot analysis of all other proteins, protein samples were separated with Novex 10-20% Tris-glycine mini gels (Life Technologies). A total of  $5 \times 10^5$  cells was loaded per lane for Chlamydomonas samples, and a total of 1.5  $\mu g$  of chlorophyll was loaded per lane for Arabidopsis samples. Proteins were then transferred to nitrocellulose membranes, blocked with 5% non-fat dry milk and blotted with specific polyclonal antibodies. The signals were detected by Supersignal West Femto Chemiluminescent substrate detection system (ThermoScientific).

*Chlamydomonas* cell fractionation, *Chlamydomonas* and *Arabidopsis* thylakoid preparations, protease protection assays and western experiments were successfully repeated three times.

**Polypeptide extraction from thylakoid.** Freshly isolated thylakoids were resuspended at 0.5 mg of chlorophyll ml<sup>-1</sup> in thylakoid resuspension buffer (0.3 M sorbitol, 2.5 mM EDTA, 5 mM MgCl<sub>2</sub>, 0.5% (wt/vol) BSA, 20 mM HEPES (pH 7.6)) containing 2 M NaBr, or 0.1 M Na<sub>2</sub>CO3, or 2 M NaSCN or no additive. After incubation on ice for 30 min, the membrane and the supernatant fraction were separated by centrifugation at 20,000*g*, 4 °C for 10 min. The membrane fractions were washed three times before being resuspended with 1× PBS buffer containing 1 mM PMSF. The supernatants were precipitated in 80% acetone and centrifuged at 20,000*g*, 4 °C for 10 min to collect pellets. The pellets were then resuspended with 1× PBS buffer containing 1 mM PMSF. The membrane and supernatant fraction were subsequently subjected to immunoblot analysis.

# Received 29 April 2016; accepted 10 August 2016; published 12 September 2016

#### References

- 1. Demmig-Adams, B. Carotenoids and photoprotection in plants: a role for the xanthophyll zeaxanthin. *Biochim. Biophys. Acta* **1020**, 1–24 (1990).
- Jahns, P., Latowski, D. & Strzalka, K. Mechanism and regulation of the violaxanthin cycle: the role of antenna proteins and membrane lipids. *Biochim. Biophys. Acta* 1787, 3–14 (2009).
- Maresca, J. A., Graham, J. E., Wu, M., Eisen, J. A. & Bryant, D. A. Identification of a fourth family of lycopene cyclases in photosynthetic bacteria. *Proc. Natl Acad. Sci. USA* 104, 11784–11789 (2007).
- 4. Niyogi, K. K. Photoprotection revisited: genetic and molecular approaches. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **50**, 333–359 (1999).
- Müller, P., Li, X.-P. & Niyogi, K. K. Non-photochemical quenching. A response to excess light energy. *Plant Physiol.* 125, 1558–1566 (2001).
- Ruban, A. V., Johnson, M. P. & Duffy, C. D. P. The photoprotective molecular switch in the photosystem II antenna. *Biochim. Biophys. Acta* 1817, 167–181 (2012).
- Niyogi, K. K. & Truong, T. B. Evolution of flexible non-photochemical quenching mechanisms that regulate light harvesting in oxygenic photosynthesis. *Curr. Opin. Plant Biol.* 16, 307–314 (2013).
- Briantais, J. M., Vernotte, C., Picaud, M. & Krause, G. H. A quantitative study of the slow decline of chlorophyll *a* fluorescence in isolated chloroplasts. *Biochim. Biophys. Acta* 548, 128–138 (1979).
- Casper-Lindley, C. & Björkman, O. Fluorescence quenching in four unicellular algae with different light-harvesting and xanthophyll-cycle pigments. *Photosynth. Res.* 56, 277–289 (1998).
- Lunch, C. K. et al. The xanthophyll cycle and NPQ in diverse desert and aquatic green algae. Photosynth. Res. 115, 139–151 (2013).
- Quaas, T. et al. Non-photochemical quenching and xanthophyll cycle activities in six green algal species suggest mechanistic differences in the process of excess energy dissipation. J. Plant Physiol. 172, 92–103 (2015).
- Baroli, I., Do, A. D., Yamane, T. & Niyogi, K. K. Zeaxanthin accumulation in the absence of a functional xanthophyll cycle protects *Chlamydomonas reinhardtii* from photooxidative stress. *Plant Cell* 15, 992–1008 (2003).
- Baroli, I. & Niyogi, K. K. Molecular genetics of xanthophyll-dependent photoprotection in green algae and plants. *Phil. Trans. R. Soc. Lond. B* 355, 1385–1394 (2000).
- Havaux, M. & Niyogi, K. K. The violaxanthin cycle protects plants from photooxidative damage by more than one mechanism. *Proc. Natl Acad. Sci. USA* 96, 8762–8767 (1999).
- Niyogi, K. K., Björkman, O. & Grossman, A. R. Chlamydomonas xanthophyll cycle mutants identified by video imaging of chlorophyll fluorescence quenching. *Plant Cell* 9, 1369–1380 (1997).

NATURE PLANTS | VOL 2 | OCTOBER 2016 | www.nature.com/natureplants

### NATURE PLANTS DOI: 10.1038/NPLANTS.2016.140

- Niyogi, K. K., Grossman, A. R. & Björkman, O. Arabidopsis mutants define a central role for the xanthophyll cycle in the regulation of photosynthetic energy conversion. *Plant Cell* 10, 1121–1134 (1998).
- Anwaruzzaman, M. *et al.* Genomic analysis of mutants affecting xanthophyll biosynthesis and regulation of photosynthetic light harvesting in *Chlamydomonas reinhardtii. Photosynth Res.* 82, 265–276 (2004).
- Yamamoto, H. Y. in *Photoprotection, Photoinhibition, Gene Regulation,* and Environment Vol. 21 (eds Demmig-Adams, B., Adams, W. W. III & Mattoo, A. K.) Ch. 1, 1–10 (Advances in Photosynthesis and Respiration Series, Springer Netherlands, 2006).
- Macko, S., Wehner, A. & Jahns, P. Comparison of violaxanthin de-epoxidation from the stroma and lumen sides of isolated thylakoid membranes from *Arabidopsis*: implications for the mechanism of de-epoxidation. *Planta* 216, 309–314 (2002).
- 20. Britton, G. Later reactions of carotenoid biosynthesis. *Pure Appl. Chem.* 47, 223–236 (1976).
- Bradbury, L. M. T. *et al.* Lycopene cyclase paralog CruP protects against reactive oxygen species in oxygenic photosynthetic organisms. *Proc. Natl Acad. Sci. USA* 109, E1888–E1897 (2012).
- 22. Leliaert, F. *et al.* Phylogeny and molecular evolution of the green algae. *Crit. Rev. Plant Sci.* **31**, 1–46 (2012).
- Kathir, P. et al. Molecular map of the Chlamydomonas reinhardtii nuclear genome. Eukaryotic Cell 2, 362–379 (2003).
- Dent, R. M. *et al.* Large-scale insertional mutagenesis of *Chlamydomonas* supports phylogenomic functional prediction of photosynthetic genes and analysis of classical acetate-requiring mutants. *Plant J.* 82, 337–351 (2015).
- Dent, R. M., Haglund, C. M., Chin, B. L., Kobayashi, M. C. & Niyogi, K. K. Functional genomics of eukaryotic photosynthesis using insertional mutagenesis of *Chlamydomonas reinhardtii. Plant Physiol.* 137, 545–556 (2005).
- Müller-Moulé, P., Conklin, P. L. & Niyogi, K. K. Ascorbate deficiency can limit violaxanthin de-epoxidase activity in vivo. *Plant Physiol.* 128, 970–977 (2002).
- Horton, R. M., Cai, Z. L., Ho, S. N. & Pease, L. R. Gene splicing by overlap extension: tailor-made genes using the polymerase chain reaction. *Biotechniques* 8, 528–535 (1990).
- Earley, K. W. et al. Gateway-compatible vectors for plant functional genomics and proteomics. Plant J. 45, 616–629 (2006).

- Zhang, X., Henriques, R., Lin, S.-S., Niu, Q.-W. & Chua, N.-H. Agrobacteriummediated transformation of *Arabidopsis thaliana* using the floral dip method. *Nat. Protoc.* 1, 641–646 (2006).
- Leonelli, L., Erickson, E., Lyska, D. & Niyogi, K. K. Transient expression in *Nicotiana benthamiana* for rapid functional analysis of genes involved in non-photochemical quenching and carotenoid biosynthesis. *Plant J.* http://dx.doi.org/10.1111/tpj.13268 (2016).
- Brooks, M. & Niyogi, K. in *Chloroplast Research in Arabidopsis* Vol. 775 (ed. Paul Jarvis, R.) Ch. 16, 299–310 (Methods in Molecular Biology Series, Humana Press, 2011).
- 32. Harris, E. H. The Chlamydomonas Sourcebook (Academic Press, 1989).
- Chua, N. H. & Bennoun, P. Thylakoid membrane polypeptides of *Chlamydomonas reinhardtii*: wild-type and mutant strains deficient in photosystem II reaction center. *Proc. Natl Acad. Sci. USA* 72, 2175–2179 (1975).
- 34. Brooks, M. D., Sylak-Glassman, E. J., Fleming, G. R. & Niyogi, K. K. A thioredoxin-like/β-propeller protein maintains the efficiency of light harvesting in *Arabidopsis. Proc. Natl Acad. Sci. USA* **110**, E2733–E2740 (2013).

### Acknowledgements

We thank J. García-Cerdán and R. Calderon for helpful discussion of *Chlamydomonas* subcellular localization. This work was supported by the US Department of Energy, Office of Science, Basic Energy Sciences, Chemical Sciences, Geosciences and Biosciences Division under field work proposal 449B. K.K.N. is an investigator of the Howard Hughes Medical Institute and the Gordon and Betty Moore Foundation (through Grant GBMF3070).

#### Author contributions

Z.L., G.P., R.M.D., Y.B., W.A., S.Y.Y. and L.L. performed research; Z.L., G.P., R.M.D. and K.K.N. designed research; Z.L., G.P. and K.K.N. analysed data and wrote the paper; all authors discussed the results and commented on the manuscript.

### Additional information

Supplementary information is available for this paper. Reprints and permissions information is available at www.nature.com/reprints. Correspondence and requests for materials should be addressed to K.K.N.

### **Competing interests**

The authors declare no competing financial interests.